arXiv:1402.0829v1 [math.FA] 4 Feb 2014

Sharp Estimates for Norms of Functions from Conjugate Classes in Metrics C and L

V. F. Babenko*

Abstract

Sharp estimates for C - and L - norms of functions that are conjugate with functions from the classes W^rH_ω of periodic functions having prescribed concave majorant of moduli of continuity, as well as sharp estimates for the differences of such functions are obtained.

Let W^rH_ω ($r = 0, 1, ...;\ W^0H_\omega = H_\omega$) be the class of 2π-periodic, r times differentiable functions $f(x)$ such that the modulus of continuity of its r-th derivative $\omega(f^{(r)}, t)$ ($f^{(0)} = f$) is less or equal to the given modulus of continuity $\omega(t)$. In addition, let $W_0^rH_\omega$ be the set of functions $f \in W^rH_\omega$ satisfying

$$\int_0^{2\pi} f(t)dt = 0.$$

For a function $f(x)$, denote by $\widetilde{f}(x)$ the function that is trigonometrically conjugate (see [1]) with $f(x)$. Let

$$\widetilde{W}^rH_\omega := \{\widetilde{f}\ :\ f \in W^rH_\omega\}.$$

In papers [2], [3] for any $r = 0, 1, ...$ and concave modulus of continuity $\omega(t)$ exact values

$$M_r(\omega)_C := \sup_{f\in W_0^rH_\omega} \|f\|_C,\ \ \Omega_r(\omega, t) := \sup_{f\in W^rH_\omega} \max_x |f(x + t) - f(x)|$$

and

$$M_r(\omega)_L := \sup_{f\in W_0^rH_\omega} \|f\|_L$$

were obtained. Estimates for the values

$$\widetilde{M}_r(\omega)_C := \sup_{f\in \widetilde{W}_0^rH_\omega} \|f\|_C,\ \ \widetilde{\Omega}_r(\omega, t) := \sup_{f\in \widetilde{W}^rH_\omega} \max_x |f(x + t) - f(x)|,\ r \neq 0,$$

as well as estimates for differences of higher order were found in [4].

We have proved the following statements.

*This paper was published in Russian in Studies in Modern Problems of Summation and Approximation of Functions and Their Applications, Collect. Sci. Works, Dnepropetrovsk, 1973, pp. 3 -5

Theorem 1. *If $\omega(t)$ is concave modulus of continuity and for some $\varepsilon > 0$*

$$\int_0^{\varepsilon} \frac{\omega(t)}{t} dt < \infty,$$

then

1.

$$\widetilde{M_0}(\omega)_C := \sup_{f \in H_\omega} \|\widetilde{f}\|_C = \frac{1}{\pi} \int_0^{\pi/2} \frac{\omega(2t)}{\sin t} dt,$$

2.

$$\widetilde{\Omega}_0(\omega, t) := \sup_{f \in H_\omega} \max_x |\widetilde{f}(x+t) - \widetilde{f}(x)| = \frac{2}{\pi} \int_0^{t/2} \frac{\sin t/2}{\cos u - \cos t/2} \omega(\rho(u) - u) du,$$

where $\rho(x)$ is defined by the equation

$$\int_0^{x} \frac{\sin t/2}{\cos u - \cos t/2} du = -\int_{\rho(x)}^{\pi} \frac{\sin t/2}{\cos u - \cos t/2} du \ \ (0 < x < t/2,\ t/2 < \rho(x) < \pi).$$

Note that proof of this Theorem essentially uses the representation (see [1, p. 573])

$$\widetilde{f}(x) = -\frac{1}{\pi} \cdot (P) \int_{-\pi}^{\pi} \frac{f(t)}{2 \tan \frac{t-x}{2}} dt$$

and Lemma 5.1 from [5].

Remark 1. *There exist concave moduli of continuity, e. g. $\omega(t) = -1/\ln t$, for which the values $\widetilde{M_0}(\omega)_C$ and $\widetilde{\Omega}_0(\omega, t)$ $(t > 0)$ are equal to ∞ .*

Remark 2. *If $E_n(f)$ is the best uniform approximation of 2π - periodic function f by trigonometric polynomials of order n, then*

$$\sup_{f \in \widetilde{W}^r H_\omega} E_0(f) = \frac{1}{2} \max_t \widetilde{\Omega}_r(\omega, t) = \frac{1}{2} \widetilde{\Omega}_r(\omega, \pi) \ \ (r = 0, 1, ...).$$

Theorem 2. *For any concave modulus of continuity the following equalities hold*

$$\widetilde{M_r}(\omega)_L := \sup_{f \in \widetilde{W}^r H_\omega} \|f\|_L = 4 \sum_{i=0}^{\infty} \frac{(-1)^{i(r+1)} b_{2i+1}}{(2i+1)^{r+1}} \ \ (r = 1, 2, ...)$$

where

$$b_m = \frac{2}{\pi} \int_0^{\pi/2} \omega(2t) \sin mt \, dt.$$

Proof of this Theorem uses analogs of Lemmas 1 and 2 from [3].

Corollary 1. *For any $r = 2, 3, ...$ the following equalities hold*

$$\sup_{f \in W^r K} \|\widetilde{f}\|_L = \frac{16K}{\pi} \sum_{i=0}^{\infty} \frac{(-1)^{i(r+1)}}{(2i+1)^{r+2}}$$

where $W^r K$ is the class of 2π - periodic functions $f(x)$ such that $f^{(r-1)}$ is absolutely continuous and almost everywhere $|f^{(r)}(x)| \leq K$.

Corollary 2. *For any $r = 2, 3, ...$ and any concave modulus of continuity $\omega(t)$ the following equalities hold*

$$\sup_{f \in \widetilde{W}^r H_\omega} \bigvee_0^{2\pi}(f) = 4 \sum_{i=0}^{\infty} \frac{(-1)^{ir} b_{2i+1}}{(2i+1)^r}.$$

Remark 3. *For every presented above extremal relation there exists a function from corresponding class that realizes supremum.*

References

[1] N. K. Bari, Trigonometric series, Moscow, Fizmatgiz, 1961

[2] N. P. Korneichuk, On extremal properties of periodic functions, Dop. AN UkrSSR, № 8 (1962), pp. 993 - 998.

[3] N. P. Korneichuk, Sharp estimates for norms of differentiable periodic functions in metric L, Mat. Zametki, v. 2, № 6, (1967), pp. 569 - 576

[4] V. P. Motornyi, About one inequality for moduli of smoothness of periodic functions, in "The first Respublic conference of yung investigators. Issue 2", Institute of Mathematics of AS of UkrSSR, Kiev, (1965), pp. 519 - 525.

[5] N. P. Korneichuk, Extremal values of functionals and the best approximations on the classes of periodic functions, Izvestiya AS USSR. Ser. Mat., v. 35, № 1 (1971), pp. 93 - 124.